

# QCD where it Matters

Ralf W. Gothe

UNIVERSITY OF  
SOUTH CAROLINA

## QCD Downunder 2017

July 10-14, 2017, Cairns in Queensland, Australia

- **$\gamma NN^*$  Vertexcouplings:** A unique exploration of baryon and quark structure?
- **Analysis and New Results:** Exclusive, quasi-free, and final state interaction!
- **Outlook:** New experiments with extended scope and kinematics!


---

# Spectroscopy

# Build your Mesons and Baryons ...

Three Generations of Matter (Fermions)

	I	II	III	
mass →	2.4 MeV	1.27 GeV	171.2 GeV	0
charge →	$\frac{2}{3}$	$\frac{2}{3}$	$\frac{2}{3}$	0
spin →	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1
name →	<b>u</b> up	<b>c</b> charm	<b>t</b> top	<b>γ</b> photon
Quarks	4.8 MeV	104 MeV	4.2 GeV	0
	$-\frac{1}{3}$	$-\frac{1}{3}$	$-\frac{1}{3}$	0
	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1
	<b>d</b> down	<b>s</b> strange	<b>b</b> bottom	<b>g</b> gluon
Leptons	<2.2 eV	<0.17 MeV	<15.5 MeV	91.2 GeV
	0	0	0	0
	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1
	<b>ν<sub>e</sub></b> electron neutrino	<b>ν<sub>μ</sub></b> muon neutrino	<b>ν<sub>τ</sub></b> tau neutrino	<b>Z<sup>0</sup></b> weak force
	0.511 MeV	105.7 MeV	1.777 GeV	80.4 GeV
	-1	-1	-1	$\pm 1$
	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1
	<b>e</b> electron	<b>μ</b> muon	<b>τ</b> tau	<b>W<sup>±</sup></b> weak force




Bosons (Forces)

# Build your Mesons and Baryons ...

Three Generations of Matter (Fermions)

	I	II	III	
mass →	2.4 MeV	1.27 GeV	171.2 GeV	0
charge →	$\frac{2}{3}$	$\frac{2}{3}$	$\frac{2}{3}$	0
spin →	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1
name →	<b>u</b> up	<b>c</b> charm	<b>t</b> top	<b>γ</b> photon
Quarks	4.8 MeV	104 MeV	4.2 GeV	0
	$-\frac{1}{3}$	$-\frac{1}{3}$	$-\frac{1}{3}$	0
	<b>d</b> down	<b>s</b> strange	<b>b</b> bottom	<b>g</b> gluon
	<2.2 eV	<0.17 MeV	<15.5 MeV	91.2 GeV
	0	0	0	0
	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1
	<b>ν<sub>e</sub></b> electron neutrino	<b>ν<sub>μ</sub></b> muon neutrino	<b>ν<sub>τ</sub></b> tau neutrino	<b>Z</b> weak force
Leptons	0.511 MeV	105.7 MeV	1.777 GeV	80.4 GeV
	-1	-1	-1	$\pm 1$
	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1
	<b>e</b> electron	<b>μ</b> muon	<b>τ</b> tau	<b>W<sup>±</sup></b> weak force




$$\mathcal{L} = \frac{1}{4g^2} G_{\mu\nu}^a G_{\mu\nu}^a + \sum_j \bar{q}_j (i\gamma^\mu D_\mu + m_j) q_j$$

where  $G_{\mu\nu}^a \equiv \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + if_{abc} A_\mu^b A_\nu^c$   
and  $D_\mu \equiv \partial_\mu + it^a A_\mu^a$   
*That's it?*

Frank Wilczek, Physics Today, August 2000


Bosons (Forces)




# N and $\Delta$ Excited Baryon States ...

Simon Capstick

➤ Orbital excitations  
(two distinct kinds in contrast to mesons)



➤ Radial excitations  
(also two kinds in contrast to mesons)




# Quark Model Classification of $N^*$

Lowest Baryon Supermultiplets  
 SU(6)xO(3) Symmetry

Dietmar Menze


Particle Data Group

- \*\*\*\*
- \*\*\*
- \*\*




# Quark Model Classification of $N^*$

BnGa energy-dependent coupled-channel PWA of CLAS  $K^+\Lambda$  and other data



# N\* Spectrum in LQCD

The strong interaction physics is encoded in the nucleon excitation spectrum that spans the degrees of freedom from meson-baryon and dressed quarks to elementary quarks and gluons.




LQCD predicts states with the same quantum numbers as CQMs with underlying  $SU(6) \times O(3)$  symmetry.

R. Edwards *et al.*,  
arXiv:1104.5152, 1201.2349



# N\* Spectrum in LQCD

The strong interaction physics is encoded in the nucleon excitation spectrum that spans the degrees of freedom from meson-baryon and dressed quarks to elementary quarks and gluons.




LQCD predicts hybrid baryon states replicating the negative parity multiplet structure.

New approved experiment on electroexcited baryon hybrids (E12-16-010).


# Quark Model Classification of $N^*$

BnGa energy-dependent coupled-channel PWA of CLAS  $K^+\Lambda$  and other data




# Transition Form Factors


# Hadron Structure with Electromagnetic Probes




- Study the structure of the nucleon spectrum in the domain where dressed quarks are the major active degree of freedom.
- Explore the formation of excited nucleon states in interactions of dressed quarks and their emergence from QCD.




# Hadron Structure with Electromagnetic Probes




- Study the structure of the nucleon spectrum in the domain where dressed quarks are the major active degree of freedom.
- Explore the formation of excited nucleon states in interactions of dressed quarks and their emergence from QCD.



# Baryon Excitations and Quasi-Elastic Scattering



hard and confined




Elastic Form Factors

Transition Form Factors

hard

soft



Deep Inelastic Scattering


S. Stein et al., PR **D22** (1975) 1884

# Structure Analysis of the Baryon


Demolition of a chimney at the "Henninger Brewery" in Frankfurt am Main, Germany, on 2 December 2006




hard and confined



hard and soft



quasi-elastic



$\gamma_{\nu} NN^*$

# Extraction




# Data-Driven Data Analyses

## Consistent Results




- Single meson production:  
Unitary Isobar Model (UIM)  
Fixed- $t$  Dispersion Relations (DR)
- Double pion production:  
Unitarized Isobar Model (JM)
- Coupled-Channel Approaches:  
EBAC  $\Rightarrow$  Argonne-Osaka  
JAW  $\Rightarrow$  Jülich-Athens-Washington  $\Rightarrow$  JüBo  
BoGa  $\Rightarrow$  Bonn-Gatchina

Int. J. Mod. Phys. E, Vol. 22, 1330015 (2013) 1-99




# New LQCD Data Analysis Approach



## Scattering processes and resonances from lattice QCD

Raul A. Briceño, Jozef J. Dudek, and Ross D. Young, arXiv:1706.06223 [hep-lat]




# New LQCD Data Analysis Approach



## Scattering processes and resonances from lattice QCD

Raul A. Briceño, Jozef J. Dudek, and Ross D. Young, arXiv:1706.06223 [hep-lat]

# Electrocouplings of $N(1520)D_{13}$ and $N(1535)S_{11}$



— Argonne Osaka / EBAC DCC MB dressing  
(absolute values)

— E. Santopinto, M. Giannini, hCQM  
PRC 86, 065202 (2012)


— S. Capstick, B.D. Keister (rCQM)  
PRD51, 3598 (1995)

■  $\pi^+\pi^-p$  2012    ▲  $\pi^+\pi^-p$  2010    ●  $N\pi$  2009


▲  $\eta p$   
CLAS/Hall-C

# Interplay between Meson-Baryon Cloud and Quark Core

N(1520)3/2-



N(1675)5/2-



..... Argonne-Osaka MB dressing (absolute values)

— E. Santopinto and M. Giannini, PRC 86 (2012) 065202


The almost direct access to

- quark core from the data on N(1520)3/2-
- meson-baryon cloud from the data on N(1675)5/2-

sheds light on the transition from the confined quark to the colorless meson-baryon structure and its dependents on the N\* quantum numbers.


# Evidence for the Onset of Precocious Scaling?

I. G. Aznauryan *et al.*, Phys. Rev. C80, 055203 (2009)




➤  $A_{1/2} \propto 1/Q^3$

➤  $A_{3/2} \propto 1/Q^5$



# Evidence for the Onset of Precocious Scaling?


Ye Tian



V. Mokeev, [userweb.jlab.org/~mokeev/resonance\\_electrocouplings/](http://userweb.jlab.org/~mokeev/resonance_electrocouplings/) (2016)

# Evidence for the Onset of Precocious Scaling?

Ye Tian







V. Mokeev, [userweb.jlab.org/~mokeev/resonance\\_electrocouplings/](http://userweb.jlab.org/~mokeev/resonance_electrocouplings/) (2016)



# N(1520)D<sub>13</sub> Helicity Asymmetry

L. Tiator

$$A_{\text{hel}} = \frac{A_{1/2}^2 - A_{3/2}^2}{A_{1/2}^2 + A_{3/2}^2}$$




○ world data

▲ PDG estimation ● ■ Nπ (UIM, DR)


# $\gamma NN^*$ Helicity Asymmetries

Ye Tian





V. Mokeev, [userweb.jlab.org/~mokeev/resonance\\_electrocouplings/](http://userweb.jlab.org/~mokeev/resonance_electrocouplings/) (2016)

# Electrocouplings of $N(1440)P_{11}$ History



- Lowest mass hybrid baryon should be  $J^P=1/2^+$  as Roper.
- In 2002 Roper  $A_{1/2}$  results were consistent with a hybrid state.

# Electrocouplings of $N(1440)P_{11}$ with CLAS




- $A_{1/2}$  has zero-crossing near  $Q^2=0.5$  and becomes dominant amplitude at high  $Q^2$ .
- Consistent with radial excitation at high  $Q^2$  and large meson-baryon coupling at small  $Q^2$ .
- Eliminates gluonic excitation ( $q^3G$ ) as a dominant contribution.

Nick Tyler closes the 1-2  $\text{GeV}^2$  gap for single pion production.

# Electrocouplings of $N(1440)P_{11}$ with CLAS

PDG 2013 update



+  $q^3g$   
 +  $q^3q\bar{q}$   
 + N-Meson  
 + ...


... all have distinctively different  $Q^2$  dependencies

- $A_{1/2}$  has zero-crossing near  $Q^2=0.5$  and becomes dominant amplitude at high  $Q^2$ .
- Consistent with radial excitation at high  $Q^2$  and large meson-baryon coupling at small  $Q^2$ .
- Eliminates gluonic excitation ( $q^3G$ ) as a dominant contribution.


Nick Tyler closes the 1-2  $\text{GeV}^2$  gap for single pion production.

# Roper Transition Form Factors in DSE Approach

$N(1440)P_{11}$




J. Segovia *et al.*, Phys. Rev. Lett. **115**, 171801



DSE Contact  $x=Q^2/m_N^2$   
 DSE Realistic  
 Inferred meson-cloud contribution  
 Anticipated complete result



Importantly, the existence of a zero in  $F_2$  is not influenced by meson-cloud effects, although its precise location is.



# Roper Transition Form Factors in DSE Approach

$N(1440)P_{11}$

J. Segovia and C.D. Roberts, arXiv:1607.04405




---

# New Experimental Results & Approaches



# Single $\pi^-$ Electroproduction off the Deuteron


Ye Tian



Exclusive  $\Rightarrow$  Spectator  $\Rightarrow$  Quasi-Free  $\Rightarrow$  FSI


# Single $\pi^-$ Electroproduction off the Deuteron

Ye Tian



# Single $\pi^-$ Electroproduction off the Deuteron


Ye Tian



Below a missing momentum of 0.2 GeV the **measured data** coincides with the resolution smeared **theoretical Fermi momentum distribution**.

# Single $\pi^-$ Electroproduction off the Deuteron


Ye Tian



Gary Hollis inclusive of the bound nucleon in the Deuteron with correction of Fermi smearing.

# Single $\pi^-$ Electroproduction off the Deuteron


Ye Tian



Below a missing momentum of 0.2 GeV the **measured data** coincides with the resolution smeared **theoretical Fermi momentum distribution**.

# Single $\pi^-$ Electroproduction off the Deuteron

Ye Tian



Momentum resolution with CLAS of the reconstructed missing momentum of the second proton.

# Single $\pi^-$ Electroproduction off the Deuteron

Ye Tian

$W = 1212 \text{ MeV}$

$\Delta W = 25 \text{ MeV}$

$Q^2 = 0.5 \text{ GeV}^2$

$\Delta Q^2 = 0.2 \text{ GeV}^2$

$\cos(\theta) = -0.7$











$\Delta \cos(\theta) = 0.2$

$\cos(\theta) = 0.7$

$\phi = 20^\circ$










$\Delta \phi = 40^\circ$

$\phi = 340^\circ$



# Single $\pi^-$ Electroproduction off the Deuteron

Ye Tian





# Single $\pi^-$ Electroproduction off the Deuteron


Ye Tian

$Q^2 = 0.5 \text{ GeV}^2$

$W = 1212 \text{ MeV}$

$Q^2 = 0.7 \text{ GeV}^2$

$Q^2 = 0.9 \text{ GeV}^2$




# Single $\pi^-$ Electroproduction off the Deuteron

Ye Tian


$Q^2 = 0.5 \text{ GeV}^2$

$W = 1212 \text{ MeV}$

$W = 1488 \text{ MeV}$




$W = 1662 \text{ MeV}$



# Single $\pi^-$ Electroproduction off the Deuteron

$$\gamma d \rightarrow \pi^- p(p)$$


Ye Tian



# Exclusive $\pi^+\pi^-$ Electroproduction off the Deuteron

Iuliia Skorodina


$P_X$  of  $ep(n) \rightarrow e'p'(n)\pi^+\pi^-$



# Exclusive $\pi^+\pi^-$ Electroproduction off the Deuteron

Iuliia Skorodina


$M_X^2$  of  $ep(n) \rightarrow e'p'(n)\pi^+X$ , all particles registered



# Exclusive $\pi^+\pi^-$ Electroproduction off the Deuteron

Iuliia Skorodina

$P_X$  of  $ep(n) \rightarrow e'p'(n)\pi^+\pi^-$




# FSI in the $p(n)\pi^+ \pi^-$ Final State

Final State Interactions depend strongly on:


Iuliia Skorodina

- invariant mass of final hadron system ( $W$ )
- scattering angles of final hadrons → FSI are topology dependent

$$M_x^2 = (P_e^\mu + P_p^\mu - P_{e'}^\mu - P_{p'}^\mu - P_{\pi^+}^\mu)^2$$



fully exclusive topology



$\pi^-$  missing topology





**blue curve** – data and **red curve** – simulation

# Effective FSI Correction

Iuliia Skorodina

$$\frac{d\sigma_{corrected}}{dW dQ^2 d\tau} = \frac{d\sigma_{not\ corrected}}{dW dQ^2 d\tau} F_{fsi}(\Delta W, \Delta Q^2)$$


$$F_{fsi}(\Delta W, \Delta Q^2) = \frac{\text{Area under green}}{\text{Area under red}}$$






# Integrated Cross Section off the Proton in Deuteron

Iuliia Skorodolina



# Comparison with Free Proton Cross Section

Iuliia Skorodina




**Black bullets** – free proton cross sections ( $e1e$  at  $E_{\text{beam}} = 2.039 \text{ GeV}$ )  
error bars show both statistical and systematical uncertainties  
G. Fedotov analysis note approved

**Red bullets** – bound proton quasi-free cross sections ( $e1e$  at  $E_{\text{beam}} = 2.039 \text{ GeV}$ )  
error bars show statistical uncertainty only

# CLAS12



# CLAS12 Baseline Equipment


- Optimized for exclusive and semi-inclusive reactions
- Large coverage in  $\theta$  and  $\phi$  angles
- Small angle capabilities
- Design operating luminosity of  $10^{35} \text{ cm}^{-2}\text{s}^{-1}$
- Particle ID up to high momenta for  $e^-/\pi$ ,  $\gamma/\pi^0$ , and  $\pi/K/p$  separation
- Good momentum and angle resolution
- Operate Polarized Target




# New Forward Time of Flight Detector for CLAS12

ToF12 Time Resolution Measurements





# Anticipated $N^*$ Electrocouplings from Combined Analyses of $N\pi/N\pi\pi$





Open circles represent projections and all other markers the available results with the 6-GeV electron beam

- Examples of **published and projected results** obtained within **60d** for three prominent excited proton states from analyses of  $N\pi$  and  $N\pi\pi$  electroproduction channels. Similar results are expected for many other resonances at higher masses, e.g.  $S_{11}(1650)$ ,  $F_{15}(1685)$ ,  $D_{33}(1700)$ ,  $P_{13}(1720)$ , ...
- The approved CLAS12 experiments E12-09-003 (NM,  $N\pi\pi$ ) and E12-06-108A (KY) are currently **the only experiments** that can provide data on  $\gamma_v NN^*$  electrocouplings for almost all well established excited proton states at the highest photon virtualities ever achieved in  $N^*$  studies up to  $Q^2$  of 12  $\text{GeV}^2$ , see <http://boson.physics.sc.edu/~gothe/research/pub/whitepaper-9-14.pdf>.

# Summary

- First high precision photo- and electroproduction data have become available and led to a new wave of significant developments in reaction and QCD-based theories.
- New high precision hadro-, photo-, and electroproduction data off the proton and the neutron will stabilize coupled channel analyses and expand the validity of reaction models, allowing us to
  - investigate and search for baryon hybrids (E12-16-010) ,
  - establish a repertoire of high precision spectroscopy parameters, and
  - measure light-quark-flavor separated electrocouplings over an extended  $Q^2$ -range, both to lower and higher  $Q^2$ , for a wide variety of  $N^*$  states (E12-16-010 A).
- Comparing these results with DSE, LQCD, LCSR, and rCQM will build further insights into
  - the strong interaction of dressed quarks and their confinement,
  - the emergence of bare quark dressing and dressed quark interactions from QCD, and
  - the QCD  $\beta$ -function and the origin of 98% of nucleon mass.
- A close collaboration of experimentalists and theorists has formed and is needed to push these goals, see Review Article *Int. J. Mod. Phys. E*, Vol. 22, 1330015 (2013) 1-99, that shall lead to a QCD theory that describes the strong interaction from current quarks to nuclei. **INT2016 & NSTAR2017.**





# 11<sup>th</sup> International Workshop on the Physics of Excited Nucleons

## N<sup>\*</sup>STAR 2017

## N<sup>\*</sup>STAR 2017

- ✓ Baryon spectrum through meson photoproduction
- ✓ Baryon resonances in experiments with hadron beams and in the  $e^+e^-$  collisions
- ✓ Baryon resonances in ion collisions and their role in cosmology
- ✓ Baryon structure through meson electroproduction, transition form factors, and time-like form factors
- ✓ Amplitude analyses and baryon parameter extraction
- ✓ Baryon spectrum and structure from first principles of QCD
- ✓ Advances in the modeling of baryon spectrum and structure
- ✓ Facilities and future projects
- ✓ Other topics related to  $N^*$  physics

August 20-23, 2017

University of South Carolina, Columbia, SC

<http://nstar2017.physics.sc.edu/>